

Osborn, H. F. — Estimating the age of the mastodon.

(1910).

0

HARVARD UNIVERSITY.



LIBRARY

OF THE

MUSEUM OF COMPARATIVE ZOOLOGY

35,154

GIFT OF

William Mc M Woodworth

March 14, 1910

MAR 14 1910

35,154

ADVANCE PRINT FROM THE PROCEEDINGS OF
The Seventh International Zoological Congress
BOSTON MEETING, AUGUST 19-24, 1907

A MEANS OF ESTIMATING THE AGE OF THE
MASTODON AND OTHER PROBOSCIDEA

HENRY FAIRFIELD OSBORN



CAMBRIDGE, MASSACHUSETTS
Sm 1910

11111
1000000000
1111111111



A MEANS OF ESTIMATING THE AGE OF THE MASTODON AND OTHER PROBOSCIDEA

(ABSTRACT)

HENRY FAIRFIELD OSBORN

THE skeleton known as the Warren Mastodon, discovered in Newburgh, N. Y., in 1845, and monographed by John Collins Warren, has been remounted recently in the American Museum of




FIG. 1.

Natural History. In repairing the tusks, the outer sheathing of the dentine was found in large part absent. The inner sheathing exposed a series of concentric constrictions and expansions which were observed to be approximately symmetrical on the two sides, as indicated by the series of + signs in Fig. 3, *A* and *B*. Secondly, it was noted that the intervals between these constrictions are broader in the middle and fore part of the tusks (corresponding with the youthful stage of growth of the tusk) and become narrower toward the base of the tusk (corresponding with the mature or adult stages of growth). Eighteen of these rings are preserved on one side and thirteen on the




FIG. 2.




FIG. 3.

other. They are faintly indicated also in the waving surface of the dentine of the tusk (Fig. 3).

On the hypothesis that these are actual annular increments of growth, the right tusk (Fig. 3, *A*) consisted of about twenty-eight segments, which allowing for the milk dentition and for the part worn off at the tip would assign to the Warren Mastodon an age of perhaps thirty years.

Similar annular constrictions are observed in the tusks of the mammoth from Alaska; and are also indicated in the tusks of the African elephant. Since the age of the Indian elephant and the rate of tusk growth is definitely known, the identification of similar concentric annular growths would be the means of testing the value of this hypothesis.

FIG. 1. Skeleton of Warren Mastodon as recently mounted in the American Museum of Natural History.

FIG. 2. Anterior view of skull of Warren Mastodon showing the true position of the tusks in the mastodon.

FIG. 3. Inferior and external views of the right and left tusks, showing the supposed annual growth segments.

A, right tusk viewed from the outside and somewhat from below.

B, left tusk, direct external or outside view.

Gaylord Bros.
Market
Syracuse, N. Y.
941 JAN 21 1908



3 2044 107 335 986

